

The $SU_{p,k}(2)$ quantum algebra with two parameters p and k [1] is defined by three generators $J_0, J_{\pm}$ with the following commutation relations

$$J_0 J_{\pm} - J_{\pm} J_0 = \pm J_{\pm} \quad (1)$$

$$J_+ J_- - \frac{p}{k} J_- J_+ = [[2 J_0]] \quad (2)$$

and the hermitian properties

$$J_0^+ = J_0, (J_{\pm})^+ = J_{\mp} \quad (3)$$

where

$$[[x]] = \frac{k^x - p^{-x}}{k - p^{-1}}. \quad (4)$$

This algebra can be obtained from a single parameter $SU_q(2)$ algebra defined by three generators $L_0, L_{\pm}$ having the commutation relations

$$L_0 L_{\pm} - L_{\pm} L_0 = \pm L_{\pm} \quad (5)$$

$$L_+ L_- - L_- L_+ = [2 L_0] \quad (6)$$

$$(L_0)^+ = L_0, (L_{\pm})^+ = L_{\mp} \quad (7)$$

where

$$[x] = \frac{q^x - q^{-x}}{q - q^{-1}}. \quad (8)$$

The substitutions

$$L_0 = J_0 \quad (9)$$

$$L_+ = \left(\frac{p}{k}\right)^{\frac{1}{2} J_0 - \frac{1}{4}} J_+ \quad (10)$$

$$L_- = J_- \left(\frac{p}{k}\right)^{\frac{1}{2} J_0 - \frac{1}{4}} \quad (11)$$

bring the defining relations (5-8) of the $SU_q(2)$ algebra (the L algebra) into the relations (1-4) satisfied by the two parameters $SU_{p,k}(2)$ algebra (the J algebra). In other words, the $SU_{p,k}(2)$ algebra is generated by the $SU_q(2)$ algebra. This fact leads to some relations between the quantities appearing in the representations of the $SU_{p,k}(2)$ and the corresponding ones in the representations of the $SU_q(2)$ algebra.

First remark: The Casimir operator C_q of the L algebra defined as

$$C_q = L_- L_+ + [L_0] [L_0 + 1] \quad (12)$$

is identical with the Casimir operator $C_{p,k}$ of the J algebra because the generators $J_{0,\pm}$ are in fact functions of $L_{0,\pm}$. It results that $L_{0,\pm}$ as well as $J_{0,\pm}$ commute with C_q which means that C_q is an invariant of the J algebra and hence

$$C_{p,k} = C_q. \quad (13)$$

Second remark: The eigenstates $|lm\rangle$ of the $(2l+1)$ dimensional irreducible representation of the L algebra are in the same time eigenstates of the J algebra as a consequence of the fact that these states are eigenstates of the Casimir operator $C_{p,k} = C_q$ and of the generator $L_0 = J_0$.

Third remark: The matrix elements of $J_{\pm}$ are equal with those of $L_{\pm}$ up to a factor which can be easily derived from the equations (9-11) and the second remark.

$$\langle l m + 1 | J_+ | l m \rangle = \left(\frac{p}{k}\right)^{-\frac{1}{2}m - \frac{1}{4}} ([l - m] [l + m + 1])^{\frac{1}{2}} \quad (14)$$

$$\langle l m - 1 | J_- | l m \rangle = \left(\frac{p}{k}\right)^{-\frac{1}{2}m + \frac{1}{4}} ([l + m] [l - m + 1])^{\frac{1}{2}}. \quad (15)$$

Fourth remark: If $J_{0\pm}(1)$ is a set of generators commuting with a second set of generators $J_{0\pm}(2)$, then one can find a third set of generators $J_{0\pm}(1, 2)$ defined as follows

$$J_0(1, 2) = J_0(1) + J_0(2) \quad (16)$$

$$J_{\pm}(1, 2) = J_{\pm}(1) p^{-J_0(2)} + k^{J_0(1)} L_{\pm}(2) \quad (17)$$

which have the commutation relations and the hermitian properties (1-4).

The corresponding generators of the L algebra defined by

$$L_0(1, 2) = L_0(1) + L_0(2) \quad (18)$$

$$L_{\pm}(1, 2) = L_{\pm}(1) q^{-L_0(2)} + q^{L_0(1)} L_{\pm}(2) \quad (19)$$

can be transformed into $J_0(1, 2)$ and $J_{\pm}(1, 2)$ by using the substitution (9-11).

Fifth remark: The Clebsch Gordan coefficients of the L algebra are identical with the Clebsch Gordan coefficients of the J algebra because the eigenstates $|l_1 l_2 lm\rangle$ are labeled by the eigenvalues of the same set of operators in both cases.

This remark remains valid for all recoupling coefficients.

Sixth remark: The $(2\lambda + 1)$ components $T_{\lambda\mu}$ of an irreducible tensor in the L algebra remain unmodified in the J algebra. The relations

$$(L_{\pm} T_{\lambda\mu} - q^{\mu} T_{\lambda\mu} L_{\pm}) q^{L_0} = ([\lambda \mp \mu] [\lambda \pm \mu + 1])^{\frac{1}{2}} T_{\lambda\mu\pm 1} \quad (20)$$

which define a tensor in the L algebra [2] are changed into

$$(J_{\pm} T_{\lambda\mu} - k^{\mu} T_{\lambda\mu} J_{\pm}) p^{J_0} = \left(\frac{p}{k}\right)^{\lambda - \mu - \frac{1}{2} \pm \frac{1}{2}} ([[\lambda \mp \mu]] [[\lambda \pm \mu + 1]])^{\frac{1}{2}} T_{\lambda\mu\pm 1} \quad (21)$$

for the J algebra.

The fact that one can go from the equation (20) to equation (21) by using the substitutions (9-11) shows that $T_{\lambda\mu}$ is the same in both algebras.

References

- [1] Schirrmacher A, Wess J and Zumino B 1991 Z. Phys. C **49** 317
 Ogievetsky O and Wess J 1991 Z. Phys. C **50** 123
 Smirnov Yu F and Wehrhahn R F 1992 J. Phys. A: Math. Gen. **25** 5563
- [2] Biedenharn L C and Tarlini M 1990 Lett. Math. Phys. **20** 271